

Lake Profile Brief


This is based on the results of Multiple Lake Threat Assessment and its Scenario Analysis. Refer to the Technical Report for details.



Lago de Yacyreta

Geographic Information

Lago de Yacyreta is a reservoir constructed on the La Plata River for hydropower production for Paraguay and Argentina. Most of the produced energy is utilized in Argentina, with a small portion going to Paraguay. Some criticized the project for an inadequate assessment of needs and environmental damage of the local ecology prior to its construction. Its flooding resulted in the relocation of an estimated 11,000 animals from 110 different species, as well as the relocation of 40,000 people. Nevertheless, the area is reported to have an abundant fauna and fishing areas. A ship lock was built on the Argentine side of the river to ease navigation, as was a fish ladder to aid in fish migration. The lake has long faced some serious environmental challenges, again becoming a subject for potential GEF consideration that would require an appropriately-established international consultative process, including an assessment of the lake's current scientific status.




TWAP Regional Designation	Southern America	Lake Basin Population (2010)	64,421,204
River Basin	La Plata	Lake Basin Population Density (2010; # km⁻²)	55.0
Riparian Countries	Argentina, Paraguay	Average Basin Precipitation (mm yr⁻¹)	1,454
Basin Area (km²)	810,470	Shoreline Length (km)	1,156
Lake Area (km²)	1,109	Human Development Index (HDI)	0.73
Lake Area:Lake Basin Ratio	0.001	International Treaties/Agreements Identifying Lake	Yes




THE MEADOWS CENTER
FOR WATER AND THE ENVIRONMENT
TEXAS STATE UNIVERSITY

Lago de Yacyreta Basin Characteristics



(a) Lago de Yacyreta basin and associated transboundary water systems



(b) Lago de Yacyreta basin land use

Lago de Yacyreta Threat Ranking

A serious lack of global-scale uniform data on the TWAP transboundary in-lake conditions required their potential threat risks be estimated on the basis of the characteristics of their drainage basins, rather than in-lake conditions. Using basin characteristics to rank transboundary lake threats precludes consideration of the unique features that can buffer their in-lake responses to basin-derived disturbances, including an integrating nature for all inputs, long water retention times, and complex, non-linear response dynamics.

The lake threat ranks were calculated with a spreadsheet-based interactive scenario analysis program, incorporating data and information about the nature and magnitude of their basin-derived stresses, and their possible impacts on the sustainability of their ecosystem services. These descriptive data for Lago de Yacyreta and the other transboundary lakes included lake and basin areas, population numbers and densities, areal extent of basin stressors on the lake, data grid size, and other components considered important from the perspective of the user of the data results. The scenario analysis program also provides a means to define the appropriate context and preconditions for interpreting the ranking results.

The Lago de Yacyreta threat ranks are expressed in terms of the Adjusted Human Water Security (Adj-HWS) threats, Reverse Biodiversity (RvBD) threats, and the Human Development Index (HDI) score, as well as combinations of these indices. However, it is emphasized that, being based on specific characteristics and assumptions regarding Lago de Yacyreta and its basin characteristics, the calculated threat scores represent only one possible set of lake threat rankings. Defining the appropriate context and preconditions for interpreting the lake rankings remains an important responsibility of those using the threat ranking results, including lake managers and decision-makers.

Table 1. Lago de Yacyreta Relative Threat Ranks, Based on Adjusted Human Water Security (Adj-HWS) and Reverse Biodiversity Threats, and Human Development Index (HDI) Score

(Estimated risks: red – highest; orange – moderately high; yellow – medium; green – moderately low; blue – low)

Adjusted Human Water Security (Adj-HWS) Threat Score	Relative Adj-HWS Threat Rank	Reverse Biodiversity (RvBD) Threat Score	Relative RvBD Threat Rank	Human Development Index (HDI) Score	Relative HDI Rank
0.75	37	0.66	19	0.73	35

It is emphasized that the Lago de Yacyreta rankings above are discussed here within the context of the management and decision-making process, rather than as strict numerical ranks. Based on its geographic, population and socioeconomic assumptions used in the scenario analysis program, the calculated Adj-HWS score for Lago de Yacyreta indicates a moderately low threat rank compared to other priority transboundary lakes.

The Reverse Biodiversity (RvBD) for Lago de Yacyreta, which is meant to describe its biodiversity sensitivity to basin-derived degradation, increases the lake threat to a moderately high threat rank,

compared to the other transboundary lakes. Management interventions directed to improving the biodiversity status must be viewed with caution, however, since we lack sufficient knowledge and experience to accurately predict the ultimate impacts of biodiversity manipulations and preservation efforts. Further, the RvBD scores indicate the relative sensitivity of a lake basin to human activities, and high threat scores *per se* do not necessarily justify management interventions. Such interventions may actually increase biodiversity degradation, noting that many developed countries have already fundamentally degraded their biodiversity because of economic development activities. Thus, activities undertaken to address the Adj-HWS threats may actually degrade the biodiversity status and resources, even if the health and socioeconomic conditions of the lake basin stakeholders are improved as a result of better conditions, thereby increasing stakeholder resource consumption.

The relative Human Development Index (HDI) places the Lago de Yacyreta basin in a moderately low threat rank in regard to its health, educational and economic conditions.

Table 2. Lago de Yacyreta Threat Ranks, Based on Multiple Ranking Criteria
(Scores for Adj-HWS, RvBD and HDI ranks are presented in Table 1; the ranks may differ in some cases because of rounding of tied threat scores; Estimated risks: red – highest; orange – moderately high; yellow – medium; green – moderately low; blue – low)

Adj-HWS Rank	HDI Rank	RvBD Rank	Sum Adj-HWS + RvBD	Relative Threat Rank	Sum Adj-HWS + HDI	Relative Threat Rank	Sum Adj-HWS + RvBD + HDI	Overall Threat Rank
33	36	20	58	32	74	38	94	34

When multiple ranking criteria are considered together in the threat rank calculations, the Adj-HWS and HDI scores considered together place Lago de Yacyreta in the lower third of the threat ranks. The relative threat is somewhat reduced when the Adj-HWS and RvBD threats are considered together. Considering all three ranking criteria together, Lago de Yacyreta exhibits a moderately low threat ranking.

Further, a series of parametric sensitivity analyses of the ranking results also was performed to determine the effects of changing the importance of specific criteria on the relative transboundary lake rankings. This analysis involved increasing or decreasing the weights applied to the threat ranks derived from multiple ranking criteria to reassess the relative impacts of the weight combinations on the threat ranks. For example, in determining the sensitivity of the Adjusted Human Water Security (Adj-HWS) and Biodiversity (BD) ranking criteria, the threat rank associated with the first was assumed to be of complete (100%) importance (i.e., rank weight of 1.0), while the other was assumed to be of no (0%) importance (i.e., rank weight of 0.0). The relative importance of the two ranking criteria was then successively changed, with weight combinations of 0.9 and 0.1, 0.8 and 0.2, etc., until the first ranking criteria (Adj-HWS) was assumed to be of no importance (rank weight of 0.0) and the second (BD) was of complete importance (rank weight of 1.0). In the case of Lago de Yacyreta, the 0.5 and 0.5 weight combinations for three cases of parametric analysis for Lago de Yacyreta resulted in respective threat rankings of 4th, 5th and 4th, respectively, among the total of 6 South American transboundary lakes in the TWAP study (see Technical Report, Section 4.3.3, pp44-51).

In essence, therefore, identifying potential management intervention needs for Lago de Yacyreta must be considered on the basis of both educated judgement and accurate representations of its situation. A

fundamental question to be addressed, therefore, is how can one decide that a given management intervention will produce the greatest benefit(s) for the greatest number of people in the Lago de Yacyreta basin? Accurate answers to such questions for Lago de Yacyreta, and other transboundary lakes, will require a case-by-case assessment approach that considers the specific lake situation and context, the anticipated improvements from specific management interventions, and its interactions with water systems to which the lake is linked.